

#M20004

GFP-Tag (7G9) Mouse mAb



- 50 μ l (25 Western mini-blot)
- 100 μ l (50 Western mini-blot)
- 200 μ l (100 Western mini-blot)

Orders ■ 400-6123-828
orders@ab-mart.com

Web ■ www.ab-mart.com.cn

BACKGROUND

The green fluorescent protein (GFP) was originally identified as a protein involved in the bioluminescence of the jellyfish *Aequorea victoria*. GFP cDNA produces a fluorescent product when expressed in prokaryotic cells, without the need for exogenous substrates or cofactors, making GFP a useful tool for monitoring gene expression and protein localization in vivo. Several GFP mutants have been developed, including EGFP, which fluoresce more intensely than the wildtype GFP and have shifted excitation maxima, making them useful for FACS and fluorescence microscopy as well as double-labeling applications. GFP is widely used in expression vectors as a fusion protein tag, allowing expression and monitoring of heterologous proteins fused to GFP.

REFERENCES

1. Prasher, D.C., et al. 1992. Primary structure of the *Aequorea victoria* green fluorescent protein. *Gene* 111: 229-233.
2. Chalfie, M., et al. 1994. Green fluorescent protein as a marker for gene expression. *Science* 263: 802-805.

SOURCE

This Abmart monoclonal antibody is produced by immunizing animals with full-length recombinant GFP.

SPECIFICITY

GFP-Tag (7G9) Mouse mAb detects GFP and GFP fusion proteins.

STORAGE

Store at -20°C . Stable for one year from the date of shipment.

REACTIVITY

All

ISOTYPE

Mouse IgG1

IMPORTANT


Use an **anti-MOUSE** secondary antibody to detect the 7G9 antibody.

RECOMMENDED ANTIBODY DILUTIONS


Western blotting	1:5000
Immunofluorescence	1:500
Immunoprecipitation	1:50
ELISA	1:2000

*** For Western blots, incubate membrane with diluted antibody in 5% w/v nonfat dry milk, 1 \times TBS, 0.05% Tween-20 at 4 $^{\circ}\text{C}$ with gentle shaking, overnight.**


APPLICATION DATA



Western blot analysis of over-expressed GFP-tagged protein in 293T cell lysate, using Abmart GFP-tag (7G9) Mouse mAb. The antibody dilutions are 1:5000 (lane 1) and 1:10000 (lane 2). Each lane was loaded with 10 μ g of cell lysate.



IP of extracts from 293T cells transfected (lane 1,2) with a GFP-tagged protein and a mock transfected (lane 3, using the same protein without the GFP tag), using Abmart GFP-Tag (7G9) Mouse mAb and probed on Western blot using the same antibody. Dilution: 1:50 (1), 1:100 (2), 1:50 (3).



IF analysis of 293T cells transfected with a GFP-tagged protein, using Abmart GFP-Tag (7G9) Mouse mAb at a 1:500 dilution.

COMPANION PRODUCTS

- #M20001 His-Tag (2A8) Mouse mAb
- #M20002 Myc-Tag (19C2) Mouse mAb
- #M20003 HA-Tag (26D11) Mouse mAb
- #M20007 GST-Tag (12G8) Mouse mAb
- #M20008 DYDDDDDK-Tag (3B9) Mouse mAb (Binds to same epitope as Sigma's Anti-FLAG[®] M2 Antibody)
- #M20012 Anti-Myc-Tag Mouse mAb (Agarose Conjugated)
- #M20013 Anti-HA-Tag Mouse mAb (Agarose Conjugated)
- #M20018 Anti-DYKDDDDK-Tag Mouse mAb (Agarose Conjugated) (Binds to same epitope as Sigma's Anti-FLAG[®] M2 Antibody)