#P37961-9 TriMethyl-Histone H3-K56 Rabbit pAb(H3K56me3)

🗆 50 µl

🗆 100 µl

BACKGROUND

Histones are basic nuclear proteins that are responsible for the nucleosome structure of the chromosomal fiber in eukaryotes. Nucleosomes consist of approximately 146 bp of DNA wrapped around a histone octamer composed of pairs of each of the four core histones (H2A, H2B, H3, and H4). The chromatin fiber is further compacted through the interaction of a linker histone, H1, with the DNA between the nucleosomes to form higher order chromatin structures. This gene is intronless and encodes a replication-dependent histone that is a member of the histone H3 family. Transcripts from this gene lack polyA tails; instead, they contain a palindromic termination element. This gene is located separately from the other H3 genes that are in the histone gene cluster on chromosome 6p22-p21.3.

Alternative Names

H3.4;H3/g;H3FT;H3t;HIST3H3;Histone H3;HIST1H3A; H3K56(me3)

SOURCE

A synthetic peptide of human Tri-Methyl-Histone H3-K56.

STORAGE

Store at -20 °C Stable far one year from the date of shipment.

REACTIVITY

Human, Mouse, Rat, Other (Wide Range)

ISOTYPE


Rabbit lgG

RECOMMENDED ANTIBODY DILIITIONS

WB 1:500-1:2000 IF/IHC/IP 1:50-1:200 ChIP 1:20-1:100

Orders	•	
Web	•	orders@ab-mart.com www.ab-mart.com.cn

APPLICATION



Western blot analysis of extracts from HeLa using H3K56me3 antibody.