#M22003 Rabbit Anti-human IgA HRP Order Order Orders@ab-mart.com Support 400-6123-828 support1@ab-mart.com Web www.abmart.cn

Description:

Monomeric IgA constitutes 5-15 % of the serum immunoglobulins whereas dimeric IgA is localized to mucosa surfaces such as saliva, gastrointestinal secretion, bronchial fluids and milk. Mucosal IgA plays a major role in host defence by neutralising infectious agents at mucosal surfaces. The production is usually local and antigen specific IgA producing B-cells can be found in regions under the lamina propria where they mature into dimeric IgA producing plasma cells. IgA deficiency is the most common immunodeficiency that may affect both serum and mucosal produced IgA.

Alternative Names: Immunoglobulin A

Reactivity: Human IgA

Mol.Wt.: ~55 kDa /~25 kDa

Storage Condition: Store at -20 °C. Stable for 12 months from date of receipt.

Application:

```
WB=1:1000-10000 ELISA=1:1000-10000 IHC-P=1:100-1000 IHC-F=1:100-1000
```

其他推荐产品

```
#M20001 His-Tag (2A8) Mouse mAb

#M20002 Myc-Tag (19C2) Mouse mAb

#M20003 HA-Tag (26D11) Mouse mAb

#M20004 GFP-Tag (7G9) Mouse mAb

#M20007 GST-Tag (12G8) Mouse mAb

#M20008 DYDDDDDK-Tag (3B9) Mouse mAb (Binds to same epitope as Sigma's Anti-
FLAG M2 Antibody)

#M20012 Anti-Myc-Tag Mouse mAb (Agarose Conjugated)

#M20013 Anti-HA-Tag Mouse mAb (Agarose Conjugated)

#M20018 Anti-DYKDDDDK-Tag Mouse Antibody (Agarose Conjugated) (Same as Sigma's Anti-FLAG M2)

#M20118 Anti-DYKDDDDK-Tag Mouse Antibody (Magnetic Beads) (Same as Sigma's Anti-FLAG M2)
```