

T58948

IL8 Antibody



Order 021-34695924
orders@ab-mart.com
Support 400-6123-828
support1@ab-mart.com
Web www.ab-mart.com.cn

100µl;50µl

Background:

IL-8 is a chemotactic factor that attracts neutrophils, basophils, and T-cells, but not monocytes. It is also involved in neutrophil activation.

Alternative Name:

Interleukin-8 (IL-8) (C-X-C motif chemokine 8) (Chemokine (C-X-C motif) ligand 8) (Emoctakin) (Granulocyte chemotactic protein 1) (GCP-1) (Monocyte-derived neutrophil chemotactic factor) (MDNCF) (Monocyte-derived neutrophil-activating peptide) (MONAP) (Neutrophil-activating protein 1) (NAP-1) (Protein 3-10C) (T-cell chemotactic factor) [Cleaved into: MDNCF-a (GCP/IL-8 protein IV) (IL8/NAP1 form I); Interleukin-8 ((Ala-IL-8)77) (GCP/IL-8 protein II) (IL-8(1-77)) (IL8/NAP1 form II) (MDNCF-b); IL-8(5-77); IL-8(6-77) ((Ser-IL-8)72) (GCP/IL-8 protein I) (IL8/NAP1 form III) (Lymphocyte-derived neutrophil-activating factor) (LYNAP) (MDNCF-c) (Neutrophil-activating factor) (NAF); IL-8(7-77) (GCP/IL-8 protein V) (IL8/NAP1 form IV); IL-8(8-77) (GCP/IL-8 protein VI) (IL8/NAP1 form V); IL-8(9-77) (GCP/IL-8 protein III) (IL8/NAP1 form VI)] CXCL8 IL8


Reactivity : Human

Predicted Reactivity : Monkey

Application : WB

Application Image

WB



Western blot analysis of IL8 expression in IL8 recombinant protein cell lysate.

Recommended Dilution : WB 1:500~1:2000

Mol Weight : 11kDa

Gene Symbol : CXCL8 IL8

Uniprot : P10145

Uniprot link:

<https://www.uniprot.org/uniprotkb/p10145>

Immunogen :

Source : Rabbit

Clonality : Monoclonal

Isotype : Rabbit

Research Area:

Angiogenesis #Chemotaxis #Inflammatory response #G protein-coupled receptor signaling pathway

Buffer:

PBS, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol

Storage:

Store at -20 °C. Stable for 12 months from date of receipt.