MA9038

Myc-tag Antibody

☐ 50ul ☐ 100 uL



Order 021-34695924

orders@ab-mart.com

Support 400-6123-828

support1@ab-mart.com

Web www.ab-mart.com.cn

Description: Mouse monoclonal antibody to Myc-tag

Alternative Names:


Source: Mouse

Uniprot:

Reactivity: N/A


Application:

WB



Western blot analysis of over-expressed Myc-tagged protein in 293T cell lysate.


IF/ICC



Immunofluorescent analysis of Myc-tag staining in 293T cells transfected with a Myc-tag protein. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary

antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark. DAPI was used to stain the cell nuclei (blue).

ΙP



Immunoprecipitation of Myc-tagged protein from HEK293T cells transfected with vector overexpressing Myc tag, using Anti-Myc-tag Antibody.

Specificity: Recognizes Myc-tag fusion proteins.

Clonality: Monoclonal

Dilution: WB (1/2000 - 1/5000), IF/IC (1/200 - 1/500), IP (1/100 - 1/200)

Gene Symbol:

Storage Condition: Store at -20°C long term.