

MA9050
VSV-G-tag Antibody



- 50ul
- 100 uL

Order 021-34695924
orders@ab-mart.com
Support 400-6123-828
support1@ab-mart.com
Web www.ab-mart.com.cn

Description : Mouse monoclonal antibody to VSV-G-tag

Alternative Names :


Source : Mouse

Uniprot :

Reactivity : N/A


Application:

WB



Western blot analysis of over-expressed VSV-G-tagged protein in 293T cell lysate.

IF/ICC



Immunofluorescent analysis of VSV-G-tag staining in 293T cells transfected with a VSV-G-tag protein. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary

antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark. DAPI was used to stain the cell nuclei (blue).

IP



Immunoprecipitation of VSV-G-tagged protein from HEK293T cells transfected with vector overexpressing VSV-G tag, using Anti-VSV-G-tag Antibody.

Specificity:

Recognizes C-terminal, internal, and N-terminal VSV-G-tag fusion proteins.

Clonality : Monoclonal

Dilution : WB (1/2000 - 1/5000), IF/IC (1/200 - 1/500), IP (1/100 - 1/200)

Gene Symbol :

Storage Condition : Store at -20°C long term.