LK51142

Biotin-Linked Glucocerebrosidase (GBA)



021-34695924 Order

orders@ab-mart.com

400-6123-828 Support

support1@ab-mart.com

Web

www.ab-mart.com.cn

UniProt: P04062


☐ 100ul;

Alternative Names:

GBA1; GCB; GLUC; Imiglucerase; Alglucerase; Beta-glucocerebrosidase; Glucosidase Beta, Acid; β-Glucosidase; D-Glucosyl-N-Acylsphingosine Glucohydrolase

Application:

WB



Reactivity: Human **Application:** WB IHC IF

Dilution:

Western blotting: 0.5-2ug/ml

Immunocytochemistry in formalin fixed cells: 5-20ug/ml

Immunohistochemistry in formalin fixed frozen section: 5-20ug/ml

Immunohistochemistry in paraffin section: 5-20ug/ml Enzyme-linked Immunosorbent Assay: 0.05-2ug/ml

Optimal working dilutions must be determined by end user.

Purification:

Antigen-specific affinity chromatography followed by Protein A affinity chromatography

Storage Condition and Buffer:

PBS, pH7.4, containing 0.02% NaN3, 50% glycerol.

Tag: Enzyme & Kinase;

Form: Liquid

Concentration: 500µg/mL