

MHY24G30

Recombinant Human Novel Coronavirus Spike glycoprotein(S) (E484K), partial (Active)



☐ 1mg;100ug;20ug

Order	021-34695924 orders@ab-mart.com
Support	400-6123-828 support1@ab-mart.com
Web	www.ab-mart.com.cn

Purity : Greater than 90% as determined by SDS-PAGE.

Endotoxin : Less than 1.0 EU/ug as determined by LAL method.

Biological Activity:

① Measured by its binding ability in a functional ELISA. Immobilized SARS-CoV-2-S1-RBD (E484K) at 2 µg/ml can bind human ACE2 (CSB-MP866317HU-B), the EC50 is 6.597-8.187 ng/ml. ② Measured by its binding ability in a functional ELISA. Immobilized SARS-CoV-2-S1-RBD (E484K) at 2 µg/ml can bind Biotin-S Antibody (CSB-RA33245D1GMY), the EC50 is 21.54-26.77 ng/ml.

Gene Names : S

Uniprot ID : P0DTC2

Organism:

Severe acute respiratory syndrome coronavirus 2 (2019-nCoV) (SARS-CoV-2)

Host : Mammalian cell

Mol.Wt. : 54.4 kDa

Expression Region : 319-541aa(E484K)

Tag Info : C-terminal mFc-tagged

Form : Lyophilized powder

Buffer:

Lyophilized from a 0.2 µm filtered PBS, 6% Trehalose, pH 7.4

Reconstitution:

We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom.


Please reconstitute protein in deionized sterile water to a concentration of 0.1-1.0 mg/mL. We recommend to add 5-50% of glycerol (final concentration) and aliquot for long-term storage at -20°C/-80°C. Our default final concentration of glycerol is 50%. Customers could use it as reference.

Notes:

Repeated freezing and thawing is not recommended. Store working aliquots at 4°C for up to one week.

Lead Time : 3-7 business days

Images:



(Tris-Glycine gel) Discontinuous SDS-PAGE (reduced) with 5% enrichment gel and 15% separation gel.

Activity



Measured by its binding ability in a functional ELISA. Immobilized SARS-CoV-2-S1-RBD (E484K) at 2 µg/ml can bind human ACE2 (CSB-MP866317HU-B), the EC₅₀ is 6.597-8.187 ng/ml.

Activity



Measured by its binding ability in a functional ELISA. Immobilized SARS-CoV-2-S1-RBD (E484K) at 2 µg/ml can bind Biotin-S Antibody (CSB-RA33245D1GMY), the EC₅₀ is 21.54-26.77 ng/ml.