

MHY24G84

Recombinant Human Novel Coronavirus Spike glycoprotein(S), partial (Active)



1mg;100ug;20ug

Order	021-34695924 orders@ab-mart.com
Support	400-6123-828 support1@ab-mart.com
Web	www.ab-mart.com.cn

Purity : Greater than 90% as determined by SDS-PAGE.

Endotoxin : Less than 1.0 EU/ug as determined by LAL method.

Biological Activity:

① Measured by its binding ability in a functional ELISA. Immobilized SARS-CoV-2-S1-RBD at 2 µg/ml can bind Biotinylated Anti-SARS-CoV-2-S Antibody (CSB-RA33245D1GMY), the EC50 of SARS-CoV-2-S1-RBD protein is 106.2-131.2 ng/ml.

② Measured by its binding ability in a functional ELISA. Immobilized human ACE2 (CSB-MP866317HU) at 2 µg/ml can bind SARS-CoV-2-S1-RBD, the EC50 of SARS-CoV-2-S1-RBD protein is 8.363-12.82 ng/ml.

Gene Names : S

Uniprot ID : P0DTC2

Organism:

Severe acute respiratory syndrome coronavirus 2 (2019-nCoV) (SARS-CoV-2)

Host : Mammalian cell

Mol.Wt. : 51.1 kDa

Expression Region : 319-541aa

Tag Info : C-terminal 6xHis-mFc-tagged

Form : Lyophilized powder

Buffer:

Lyophilized from a 0.2 µm filtered 20 mM Tris-HCl, 0.5 M NaCl, 6% Trehalose, pH 8.0

Reconstitution:


We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Please reconstitute protein in deionized sterile water to a concentration of 0.1-1.0 mg/mL. We recommend to add 5-50% of glycerol (final concentration) and aliquot for long-term storage at -20°C/-80°C. Our default final concentration of glycerol is 50%. Customers could use it as reference.

Notes:

Repeated freezing and thawing is not recommended. Store working aliquots at 4°C for up to one week.

Lead Time : 3-7 business days

Images:



(Tris-Glycine gel) Discontinuous SDS-PAGE (reduced) with 5% enrichment gel and 15% separation gel.

Predicted band size: 51.1 kDa

Observed band size: 66 kDa due to glycosylation



Measured by its binding ability in a functional ELISA. Immobilized SARS-CoV-2-S1-RBD at 2 µg/ml can bind Biotinylated Anti-SARS-CoV-2-S Antibody (CSB-RA33245D1GMY), the EC₅₀ of SARS-CoV-2-S1-RBD protein is 106.2-131.2 ng/ml.



Measured by its binding ability in a functional ELISA. Immobilized human ACE2 (CSB-MP866317HU) at 2 µg/ml can bind SARS-CoV-2-S1-RBD, the EC₅₀ of SARS-CoV-2-S1-RBD protein is 8.363-12.82 ng/ml.