MN50074

c-Myc Tag mAb



Order 021-34695924

orders@ab-mart.com

Support 400-6123-828

support1@ab-mart.com

Web www.ab-mart.com.cn

Background:

☐ 50ul;100ul;

Myc tag is a polypeptide protein tag derived from the c-myc gene product that can be added to a protein using recombinant DNA technology. It can be used for affinity chromatography, and then used to separate recombinant, overexpressed protein from wild type protein expressed by the host organism. It can also be used in the isolation of protein complexes with multiple subunits.


Alternative Names:

c-Myc; Myc tag; Myc protein; Myc epitope tag; Myc2

Source : Mouse

Application

WB



Western blot analysis of Myctag in CHOK1, 11654myc, 02103flag lysates using Myctag antibody.

Clonality: Monoclonal Antibody

Isotype: IgG1

Immunogen: Synthetic peptide EQKLISEEDL coupled to KLH.

Reactivity: Transfected

Dilution: WB: 1/1000-1/10000

Molecular Weight: Refer to figures

Purification: Affinity Purified

Form : Liquid

Concentration: 1 mg/ml

Buffer:

Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide, pH 7.3.

Storage:

Store at 4°C short term. Store at -20°C long term. Avoid freeze/thaw cycles.