

MN50178

HA Tag mAb

Abmart

Order 021-34695924
orders@ab-mart.com
Support 400-6123-828
support1@ab-mart.com
Web www.ab-mart.com.cn

□ 50ul;100ul;

Background:

The HA tag is derived from the human influenza hemagglutinin corresponding to amino acids 98-106. It has been extensively used as a general epitope tag in expression vectors. This tag facilitates the detection, isolation, and purification of the proteins.


Alternative Names:

HA epitope tag; HA1; HA2; hemagglutinin; Hemagglutinin HA1 chain; Hemagglutinin HA2 chain

Source : Mouse

Application

WB



Western blot analysis of HA-tag in CHO-K1, 09603HA, 00222PEGFP lysates using HA-tag antibody.

Clonality : Monoclonal Antibody

Isotype : IgG1

Immunogen : Synthetic Peptide of HA-Tag

Reactivity : Transfected

Dilution : WB: 1/1000-1/10000

Molecular Weight : Refer to figures

Purification : Affinity Purified

Form : Liquid

Concentration : 1 mg/ml

Buffer:

Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide, pH 7.3.

Storage:

Store at 4°C short term. Store at -20°C long term. Avoid freeze/thaw cycles.