

RMCH051

Recombinant CD27L (CD27 ligand), Human , AF

 5ug;20ug;100ug;500ug;

Order	021-34695924 orders@ab-mart.com
Support	400-6123-828 support1@ab-mart.com
Web	www.ab-mart.com.cn

Synonym/Alternative name(s):

soluble CD27 Ligand, sCD27 Ligand, TNFSF7, CD70

Activity:

Measure by its ability to induce IL-8 secretion in human PBMCs.

The ED50 for this effect is <0.6 ng/mL.

Protein Description:

CD27 Ligand, which belongs to the TNF superfamily, is a type II transmembrane protein. Like NK cells, CD27L is also located in activated T and B lymphocytes. Along with NK enhancement, CD27L and its receptor regulate the immune response by raising expansion and differentiation of T cell. CD27 signaling can act like a co-stimulatory effector to sustain survival of the CD8+ T cells, mainly by increasing expression of the IL-2 gene.

Protein Accession : P32970.2**Gene ID :** 970**Expression Sequence:**

QRFAQAQQQLPLESLGWDVAELQLNHTGPOQDPRLYWQGGPALGRSFLHGPPELDKQQLRIHRDGIYMVHIQVTLAICSSTTASRHHP
TTLAVGICSPASRSISLLRSLFHQGCTIASQRLTPLARGDTLCTNLGTLLPSRNTDETFFGVQWVRP with polyhistidine tag at the N-terminus

Fusion tag : His-tag?at?the?N-terminus**Species :** Human**Reactivity :** Human**Expression Host :** Escherichia coli**Source :** E. coli**Purity/method:**

>98% as determined by SDS-PAGE. Ni-NTA chromatography

Endotoxin level:

<0.01 EU per 1 µg of the protein by the LAL method.

Calculated Molecular Weight : 18.08 kDa**Formulation:**

The protein was lyophilized from a solution containing 1X PBS, pH 8.0.

Reconstitution:

It is recommended to reconstitute the lyophilized protein in sterile H2O to a concentration not less than 100 µg/mL and incubate the stock solution for at least 20 min to ensure sufficient re-dissolved.

Shipping : Blue Ice**Stability and Storage:**

Lyophilized protein should be stored at -20°C for 1 year.

Upon reconstitution, store at 2°C to 8°C for up to 1 week. Further dilute in a buffer containing a carrier protein or

stabilizer (e.g. 0.1%

BSA, 10%FBS, 5%HSA or 5% trehalose solution), protein aliquots should be stored at -20°C or -80°C for 3-6 months.

Category : Cytokines

Application : Cell culture, Elisa

Image:



SDS- PAGE analysis of recombinant human CD27L