

RMCH103

Recombinant FGF-13 (Fibroblast growth factor-13), Human, AF



5ug;20ug;100ug;500ug;

Order	021-34695924 orders@ab-mart.com
Support	400-6123-828 support1@ab-mart.com
Web	www.ab-mart.com.cn

Synonym/Alternative name(s) : FHF-2, FHF2, LINC00889

Activity:

Measure by its ability to induce 3T3 cells proliferation.

The ED50 for this effect is <160 ng/mL.

Protein Description:

Fibroblast growth factor 13 (FGF13) is a new member of the fibroblast growth factor (FGF) family. FGF family members possess broad mitogenic and cell survival activities and are involved in a variety of biological processes, including embryonic development, cell growth, morphogenesis, tissue repair, tumor growth, and invasion. FGF-13 plays an important role in the regulation of embryonic development and as a signaling molecule in the induction and patterning of the embryonic brain.

Protein Accession : Q92913.1

Gene ID : 2258

Expression Sequence:

MAAAIASSLIRQKRQAREREKSNACKCVSSPSKGTSCDKNKLNVFSRVKLFSGSKRRRRRPEPQLKGIVTKLYSRQGYHLQLQADGTID
GTKDEDSTYTLFNLIPVGLRVVAIQGVQTKLYLAMNSEGYLTSELFTEPECKFKESVFNENYVYSSMIYRQQQSGRGWYLGLNKEGEIM
KGNHVKKNKPAAHFLPKPLKVAMYKEPSLHDLTEFSRSGSGTPTKRSVSGVLNNGGKSMHNEST with polyhistidine tag at the C-terminus.

Fusion tag : His-tag?at?the?C-terminus

Species : Human

Reactivity : Human

Expression Host : Escherichia coli

Source : E. coli

Purity/method:

>98% as determined by SDS-PAGE. Ni-NTA chromatography

Endotoxin level:

<0.1 EU per 1 µg of the protein by the LAL method.

Calculated Molecular Weight : 21.23 kDa

Formulation:

The protein was lyophilized from a solution containing 1X PBS, pH 7.4.

Reconstitution:

It is recommended to reconstitute the lyophilized protein in sterile H₂O to a concentration not less than 100 µg/mL and incubate the stock solution for at least 20 min to ensure sufficient re-dissolved.

Shipping : Blue Ice

Stability and Storage:

Lyophilized protein should be stored at -20°C for 1 year.

Upon reconstitution, store at 2°C to 8°C for up to 1 week. Further dilute in a buffer containing a carrier protein or


stabilizer (e.g. 0.1%

BSA, 10%FBS, 5%HSA or 5% trehalose solution), protein aliquots should be stored at -20°C or -80°C for 3-6 months.

Category : Cytokines

Application : Cell culture, Elisa

Image:



SDS- PAGE analysis of recombinant human FGF-13