

RMCH106**Recombinant FGF-17 (Fibroblast growth factor-17), Human, AF** 5ug;20ug;100ug;500ug;

Order	021-34695924 orders@ab-mart.com
Support	400-6123-828 support1@ab-mart.com
Web	www.ab-mart.com.cn

Synonym/Alternative name(s) : FGFH**Activity:**

Measure by its ability to induce 3T3 cells proliferation.

The ED50 for this effect is <5 ng/mL.

The specific activity of recombinant human FGF-17 is > 2 x 10⁵ IU/mg.

Protein Description:

FGF-17, also known as FGF13, is a member of the fibroblast growth factor (FGF) family. FGF family members possess broad mitogenic and cell survival activities, and are involved in a variety of biological processes, including embryonic development, cell growth, morphogenesis, tissue repair, tumor growth and invasion. FGF-13 plays an important role in the regulation of embryonic development and as signaling molecule in the induction and patterning of the embryonic brain.

Protein Accession : O60258.1**Gene ID :** 8822**Expression Sequence:**

TQGENHPSPNFNQYVRDQGAMTDQLSRRQIREYQLYSRTSGKHVQVTGRRISATAEDGNKFAKLIVETDTFGSRVRIKGAESEKYICMN
KRGKLGKPSGKSKDCVFTEIVLENNYAFQNRHEGWFMFAFTRQGRPRQASRSRQNRQEAHFYKRLYQGGQLPFPNHAEKQKQFEFV
GSAPTRTRTKRTRRPQPLT with polyhistidine tag at the N-terminus.

Fusion tag : His-tag?at?the?N-terminus**Species :** Human**Reactivity :** Human**Expression Host :** Escherichia coli**Source :** E. coli**Purity/method:**

>98% as determined by SDS-PAGE. Ni-NTA chromatography

Endotoxin level:

<0.1 EU per 1 µg of the protein by the LAL method.

Calculated Molecular Weight : 23.32 kDa**Formulation:**

The protein was lyophilized from a solution containing 1X PBS, pH 8.0.

Reconstitution:

It is recommended to reconstitute the lyophilized protein in sterile H₂O to a concentration not less than 100 µg/mL and incubate the stock solution for at least 20 min to ensure sufficient re-dissolved.

Shipping : Blue Ice**Stability and Storage:**

Lyophilized protein should be stored at -20°C for 1 year.

Upon reconstitution, store at 2°C to 8°C for up to 1 week. Further dilute in a buffer containing a carrier protein or


stabilizer (e.g. 0.1%

BSA, 10%FBS, 5%HSA or 5% trehalose solution), protein aliquots should be stored at -20°C or -80°C for 3-6 months.

Category : Cytokines

Application : Cell culture, Elisa

Image:



SDS- PAGE analysis of recombinant human FGF-17