

Order 021-34695924

orders@ab-mart.com

400-6123-828 Support

support1@ab-mart.com

Web www.ab-mart.com.cn

Synonym/Alternative name(s): GRO-α: MGSAα, NAP-3, GRO1

Activity:

Measure by its ability to chemoattract BaF3 cells transfected with human CXCR2.

The ED50 for this effect is <3 ng/mL.

☐ 5ug;20ug;100ug;500ug;

Protein Description:

CXCL1/GROa act as a growth factor for melanoma cells. It is also a chemotaxin for neutrophils. Like other alpha chemokines, this protein is potent neutrophil attractant and activator and is also affect basophils. Clinical research show that CXCL1 protein may be a therapeutic target as well as a diagnostic marker in ovarian cancer.

Protein Accession: P09341

Gene ID: 2919

Expression Sequence:

ASVATELRCQCLQTLQGIHPKNIQSVNVKSPGPHCAQTEVIATLKNGRKACLNPASPIVKKIIEKMLNSDKSN with polyhistidine tag at the N-terminus.

Fusion tag: His-tag?at?the?N-terminus

Species: Human Reactivity: Human

Expression Host: Escherichia coli

Source: E. coli Purity/method:

>98% as determined by SDS-PAGE. Ni-NTA chromatography

Endotoxin level:

<0.1 EU per 1 µg of the protein by the LAL method.

Calculated Molecular Weight: 8.67 kDa

Formulation:

The protein was lyophilized from a solution containing 1X PBS, pH 7.4.

Reconstitution:

It is recommended to reconstitute the lyophilized protein in sterile H2O to a concentration not less than 100 μg/mL and incubate the stock solution for at least 20 min to ensure sufficient re-dissolved.

Shipping: Blue Ice

Stability and Storage:

Lyophilized protein should be stored at -20°C for 1 year.

Upon reconstitution, store at 2°C to 8°C for up to 1 week. Further dilute in a buffer containing a carrier protein or stabilizer (e.g. 0.1%

BSA, 10%FBS, 5%HSA or 5% trehalose solution), protein aliquots should be stored at -20°C or -80°C for 3-6 months.

Category: Cytokines

Application: Cell culture, Elisa

Image:



SDS- PAGE analysis of recombinant human CXCL1