

RMCH178**Recombinant HMGB2, Human (mammalian cell expression), AF** 5ug;20ug;100ug;500ug;

Order	021-34695924 orders@ab-mart.com
Support	400-6123-828 support1@ab-mart.com
Web	www.ab-mart.com.cn

Synonym/Alternative name(s) : HMG2**Activity :** N/A**Protein Description:**

HMGB2 is a member of the non-histone chromosomal high-mobility group protein family, which are chromatin-associated and widely expressed in the nucleus of higher eukaryotic cells. HMGB2 can assist cooperative interactions between cis-acting proteins by promoting DNA flexibility through bending DNA to DNA circles. In addition, HMGB2 participates in the final ligation step in DNA end-joining processes of DNA double-strand breaks repair and V(D)J recombination.

Protein Accession : P26583.2**Gene ID :** 3148**Expression Sequence:**

MGKGDPNKPRGKMSSYAFVQTCREEHKKKHPDSSVNFSAEFSKKCSERWKTMSAKEKSKFEDMAKSDKARYDREMKNYVPPKGDKK
GKKKDPNAPKRPPSAFFLFCSEHRPKIKSEHPGLSIGDTAKKLGEMWSEQSAKDKQPYEQKAAKLEKEYEKDIAAYRAKKGSEAGKKGP
GRPTGSKKKNEPEDEEEEEDEDEEEEEDEDEE with polyhistidine-SUMO tag at the N-terminus

Fusion tag : His-SUMO tag at the N-terminus**Species :** Human**Reactivity :** Human**Expression Host :** HEK293**Source :** HEK293**Purity/method:**

>98% as determined by SDS-PAGE. Ni-NTA chromatography

Endotoxin level:

<0.1 EU per 1 µg of the protein by the LAL method.

Calculated Molecular Weight : 35.57 kDa**Formulation:**

The protein was lyophilized from a solution containing 1X PBS, pH 7.4.

Reconstitution:

It is recommended to reconstitute the lyophilized protein in sterile H₂O to a concentration not less than 100 µg/mL and incubate the stock solution for at least 20 min to ensure sufficient re-dissolved.

Shipping : Blue Ice**Stability and Storage:**

Lyophilized protein should be stored at -20°C for 1 year.


Upon reconstitution, store at 2°C to 8°C for up to 1 week. Further dilute in a buffer containing a carrier protein or stabilizer (e.g. 0.1%

BSA, 10%FBS, 5%HSA or 5% trehalose solution), protein aliquots should be stored at -20°C or -80°C for 3-6 months.

Category : Cytokines

Application : Cell culture, Elisa

Image:



SDS- PAGE analysis of recombinant human HMGB2