RMCM028

Recombinant IL-31 (Interleukin-31), Mouse, AF



Order 021-34695924

orders@ab-mart.com

400-6123-828 Support

support1@ab-mart.com

Web www.ab-mart.com.cn

Synonym/Alternative name(s): 1700013B14Rik

Activity: N/A

Protein Description:

☐ 5ug;20ug;100ug;500ug;

Interleukin-31 (IL-31) s an inflammatory cytokine that helps trigger cell-mediated immunity against pathogens. It has also been identified as a major player in a number of chronic inflammatory diseases, including atopic dermatitis. It is produced by a variety of cells, namely type 2 helper (TH2) T-cells. IL-31 sends signals through a receptor complex made of IL-31RA and oncostatin M receptor β (OSMR β) expressed in immune and epithelial cells. These signals activate three pathways: ERK1/2 MAP kinase, PI3K/AKT, and JAK1/2 signaling pathways.

Protein Accession: Q6EAL8.1

Gene ID: 76399

Expression Sequence:

MTCSLSFGAPISKEDLRTTIDLLKQESQDLYNNYSIKQASGMSADESIQLPCFSLDREALTNISVIIAHLEKVKVLSENTVDTSWVIRWLTNI SCFNPLNLNISVPGNTDESYDCKVFVLTVLKQFSNCMAELQAKDNTTC with polyhistidine tag at the C-terminus.

Fusion tag: His-tag?at?the?C-terminus

Species: Mouse **Reactivity:** Mouse

Expression Host: Escherichia coli

Source: E. coli Purity/method:

>98% as determined by SDS-PAGE. Ni-NTA chromatography

Endotoxin level:

<0.1 EU per 1 μg of the protein by the LAL method. Calculated Molecular Weight: 16.56 kDa

Formulation:

The protein was lyophilized from a solution containing 1X PBS, pH 7.4.

Reconstitution:

It is recommended to reconstitute the lyophilized protein in sterile H2O to a concentration not less than 100 μg/mL and incubate the stock solution for at least 20 min to ensure sufficient re-dissolved.

Shipping: Blue Ice

Stability and Storage:

Lyophilized protein should be stored at -20°C for 1 year.


Upon reconstitution, store at 2°C to 8°C for up to 1 week. Further dilute in a buffer containing a carrier protein or stabilizer (e.g. 0.1%

BSA, 10%FBS, 5%HSA or 5% trehalose solution), protein aliquots should be stored at -20°C or -80°C for 3-6 months.

Category: Cytokines

Application: Cell culture, Elisa

Image:



SDS- PAGE analysis of recombinant mouse IL-31